



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

BB

Urkunde Certificate Certificat

Es wird hiermit bescheinigt, daß für die in der beigefügten Patentschrift beschriebene Erfindung ein europäisches Patent für die in der Patentschrift bezeichneten Vertragsstaaten erteilt worden ist.

It is hereby certified that a European patent has been granted in respect of the invention described in the annexed patent specification for the Contracting States designated in the specification.

Il est certifié qu'un brevet européen a été délivré pour l'invention décrite dans le fascicule de brevet ci-joint, pour les Etats contractants désignés dans le fascicule de brevet.

Europäisches Patent Nr.

European Patent No.

Brevet européen n°

0462277

Patentinhaber

Proprietor of the Patent

Titulaire du brevet


SNOW BRAND MILK PRODUCTS CO., LTD.
6-1-1, Naebo-cho Higashi-ku
Sapporo-shi Hokkaido 065/JP

München, den
Munich,
Fait à Munich, le

07.06.95


Paul Braendli

Präsident des Europäischen Patentamts
President of the European Patent Office





Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number:

0 462 277 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication of patent specification: 07.06.95 (51) Int. Cl.⁶: C07K 2/00, C07K 14/47

(21) Application number: 90904429.9

(22) Date of filing: 09.03.90

(86) International application number:
PCT/JP90/00314

(87) International publication number:
WO 90/10651 (20.09.90 90/22)

The file contains technical information submitted
after the application was filed and not included in
this specification

(54) GLYCOPROTEIN OF HUMAN ORIGIN, PHYSIOLOGICALLY ACTIVE FACTOR COMPRISING THE SAME,
AND PHARMACEUTICAL PREPARATION CONTAINING THE SAME AS ACTIVE INGREDIENT.

(30) Priority: 10.03.89 JP 58631/89
16.01.90 JP 6692/90

(43) Date of publication of application:
27.12.91 Bulletin 91/52

(45) Publication of the grant of the patent:
07.06.95 Bulletin 95/23

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB IT LI LU NL SE

(56) References cited:
JP-A- 0 110 998
JP-A-58 148 826

(73) Proprietor: SNOW BRAND MILK PRODUCTS
CO., LTD.
6-1-1, Naebo-cho
Higashi-ku
Sapporo-shi
Hokkaido 065 (JP)

(72) Inventor: HIGASHIO, Kanji
1769-10, Yamada
Kawagoe-shi
Saitama 350 (JP)
Inventor: MITSUDA, Shinjiro 3-3-8, Sakuradai
Hasuda-shi
Saitama 349-01 (JP)
Inventor: SHIMA, Nobuyuki 801-1-306,
Shikatebukuro
Urawa-shi
Saitama 336 (JP)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

BIOCHEMICAL AND BIOPHYSICAL RE-
SEARCH COMMUNICATIONS. vol. 163, no. 2,
15 September 1989, DULUTH, MINNESOTA US
pages 967 - 973; KEIJI MIYAZAWA ET
AL: "MOLECULAR CLONING AND SEQUENCE
ANALYSIS OF CDNA FOR HUMAN
HEPATOCYTE GROWTH FACTOR"

NATURE. vol. 342, 23 November 1989, LON-
DON GB pages 440 - 443;
TOSHIKAZUNAKAMURA ET AL: 'MOLECULAR
CLONING AND EXPRESSION OF HUMAN
HEPATOCYTE GROWTH FACTOR'

PROCEEDINGS OF THE NATIONAL ACADEMY
OF SCIENCES OF USA. vol. 88, January
1991, WASHINGTON US pages 415 - 419; JEF-
FREY S. RUBIN ET AL: 'A BROAD-SPECTRUM
HUMAN LUNG FIBROBLAST-DERIVED
MITOGEN IS A VARIANT OF HEPATOCYTE
GROWTH FACTOR'

NATURE. vol. 327, 21 May 1987, LONDON GB
pages 239 - 242; MICHAEL STOKER ET AL:
'SCATTER FACTOR IS A FIBROBLAST-DE-
RIVED MODULATOR OF EPITHELIAL CELL
MOBILITY.'

Proc Nat Acad Sci USA 86 (1989) 5844-48

Nature 346 (1990) 228

Inventor: ITAGAKI, Yasuharu 2-5-18-302,
Takinohara
Utsunomiya-shi
Tochigi 320 (JP)
Inventor: NAGAO, Masaya 578-15-3-4,
Ishibashi
Oaza
Ishibashicho
Shimotsugagun
Tochigi 329-05 (JP)

⑦ Representative: Davies, Jonathan Mark et al
Reddie & Grose
16 Theobalds Road
London WC1X 8PL (GB)

Description

A human-derived glycoprotein, biologically active factor which includes glycoprotein, and pharmaceutical product which comprises biologically active factor as active component.

Field of the Invention

The present invention relates to a glycoprotein obtained from the culture broth of the human-derived fibroblasts, biologically active factor which includes glycoprotein, and pharmaceutical product which comprises biologically active factor as active component.

The glycoprotein in the present invention, which shows cytotoxic activity to various tumor cell lines but not to normal cells, is a new tumor cytotoxic factor, leukemic cell differentiation inducing factor, cellular immunology enhancing factor, vascular endothelial cell growth factor and hepatocyte growth factor. This material is useful as an anti-tumor drug, an anti-leukemia drug, a cellular immunology enhancing drug, a wound healing drug and a liver regenerating drug, etc. or a biochemical or pharmacological reagent.

Description of the Related Art

β -Interferon is a representative factor as the biologically active factor, for example, the tumor cytotoxic factor which is produced by human-derived fibroblasts. This is a glycoprotein which is secreted by the fibroblasts, when after the culture, the cells are harvested and stimulated by poly I-polyC or sendai viruses. It has been clarified that the protein has various physiological activities in addition to its anti-virus or anti-tumor action. A fibroblast-derived tumor cytotoxic glycoprotein called as CBF is represented in DE 3306297. A tumor growth inhibitory factor (INF) with a molecular weight of 35,000 to 45,000 which is purified from the culture broth of human tissue-derived fibroblasts is disclosed in EP 159276. Also, a tumor necrosis factor-like material which is purified from the culture broth of fibroblasts, a fibroblast-derived necrosis factor, FNF, and a biologically active material with the cytotoxic activity, which is produced by animal fibroblast cells and has a molecular weight of 40,000 to 50,000 and an isoelectric point of 5.0 ± 0.5 , are disclosed in Japanese Laid Open No. 61-56131, Japanese Laid Open No. 61-1872, and Japanese Laid Open No. 62-103021, respectively. Furthermore, all amino acid sequence and cDNA sequence coding for the amino acid sequence of a tumor cytotoxic factor, which is obtained from the culture broth of human-derived fibroblasts, with a molecular weight of $36,000 \pm 1,000$ and an isoelectric point more than 10.5, are disclosed in Japanese Laid Open No. 64-10998.

Summary of the Invention

The present inventors have investigated about a biologically active material which is contained in the culture broth of human-derived fibroblasts and have found a glycoprotein with various biological activities, which is different from the materials reported previously in points of the molecular weight and the isoelectric point etc.

Therefore, the main aim of the present invention is to offer the new glycoprotein, biologically active factor which includes said glycoprotein, and pharmaceutical product which comprises said biologically active factor.

According to the present invention there is provided a glycoprotein which (i) is obtainable from the culture broth of human derived fibroblasts, (ii) has amino acid sequence shown in Fig. 15 and (iii) has the following physicochemical properties;

a. Molecular weight; On SDS gel electrophoresis, $78,000 \pm 2,000$ or $74,000 \pm 2,000$ under the nonreduced conditions and a common band A with $52,000 \pm 2,000$ and band B with $30,000 \pm 2,000$ or band C with $26,000 \pm 2,000$ under the reduced conditions.

b. Isoelectric point; 7.4 to 8.6.

c. Heat stability; Stable in the heating at 60°C for 10min.

d. pH stability; Stable in the range of pH6 to 9.

e. Carbohydrate chain; Adsorbed to a Concanavalin A (Con A)-Sepharose (Reg. TM) column.

f. Biological activity; Inhibit the growth of KB cells, HeLa cells, and L-929 cells but not IMR-90 cells.

g. Reactivity to antibodies; The cytotoxic activity is not neutralized by anti-TNF antibody, anti-lymphotoxin antibody, and anti-interferon- β antibody.

Furthermore, the suitable lot of TCF-II in the present invention has the following properties in addition to the above physicochemical characteristics (a-g):

h.N-terminal amino acid sequence: The above mentioned band B and band C are subchains of band A, respectively. N-terminus of band A is blocked. Band B and band C has common N-terminal amino acid sequence as follows;

Val-Val-Asn-Gly-Ile-Pro-Thr-

or

Val-Val-Asn-Gly-Ile-Pro-Thr-X-Thr-Asn-Ile-Gly-X-Met-

Val-Ser-Leu-

X means an unidentified amino acid.

i. Amino acid composition; When it is hydrolyzed with HCl, it exhibits the following amino acid composition.

A.A	nmol	mol%
Asp	10.375	12.97
Glu	7.750	9.69
Ser	5.000	6.25
Gly	7.250	9.06
His	3.000	3.75
Arg	5.375	6.72
Thr	5.125	6.41
Ala	2.625	3.28
Pro	5.625	7.03
Tyr	3.875	4.84
Val	4.125	5.16
Met	1.875	2.34
Cys	ND	-
Ile	5.000	6.25
Leu	4.875	6.09
Phe	2.250	2.81
Trp	ND	-
Lys	5.875	7.34
total	80.000	100(99.99)

Furthermore, all the primary sequence of TCF-II in the present invention was deduced from its complementary DNA (cDNA). TCF-II cDNA was cloned by screening cDNA library which was prepared by using mRNA purified from total RNA which was extracted from human embryonic fibroblast cells (IMR-90), according to the following method.

(1) Extraction of poly(A)⁺RNA from IMR-90 cells

Total RNA was prepared by guanidine thiocyanate -cesium chloride method (Biochemistry 18 5294-5299(1979)) from 2×10^8 IMR-90 cells which were cultured in the Dulbecco's modified eagle (DME) medium containing 5% of new born calf serum (NBCS). The IMR-90 cells were suspended in 28ml of 6M guanidine thiocyanate containing 5mM sodium citrate, 0.5% Sarcoseal and 0.1M β -mercaptoethanol, and were homogenized. 5.7M cesium chloride solution, 4ml containing 0.1M EDTA was put into polyallomer centrifuge tubes. The homogenized solution, 7ml was loaded on the cesium chloride solution and then centrifuged at 35,000 rpm, 20°C for 16 hours, using 40TI rotor of Beckman centrifugator. After centrifugation, the pellets were washed twice with 95% ethanol and dissolved in 200 μ l of 10mM Tris HCl buffer (pH7.5) solution containing 1mM EDTA by heating at 65°C for 5 min, designated as total RNA solution. Poly (A)⁺RNA was purified from the total RNA by the method of oligo (dT)cellulose-column chromatography. The total RNA solution was loaded on the oligo (dT) cellulose-column which was equilibrated with 10mM Tris HCl buffer(pH7.4) solution containing 1mM EDTA, 0.5M NaCl and 0.05% SDS. The adsorbed fraction was eluted with 10mM Tris HCl buffer, pH7.4, containing 1mM EDTA and 0.05% SDS, and designated as poly(A)⁺ RNA solution.

(2) Synthesis of cDNA library

Double strand cDNA was synthesized by using poly (A) ⁺RNA from (1) as a template and by using cDNA synthesis kit (Pharmacia Co. Ltd), and was attached EcoR I adaptor. The method of synthesis was performed according to the protocol of Pharmacia Co.Ltd, except addition of reverse transcriptase (40 units/reaction mixture, Life Science Co. Ltd) derived from non-sphere disease virus of avian bone marrow at the synthesis of single strand DNA.

(3) Preparation of cDNA library

The cDNA obtained from (2) was inserted in EcoR I arm (Promega Co. Ltd) of phage vector λ gt10. 3.3 μ g of cDNA synthesized from poly(A)⁺RNA was dissolved in 150 μ l of column buffer, 66 mM Tris-HCl buffer (pH7.6) containing 1mM spermidine, 10mM magnesium chloride, 15 mM dithiothreitol and bovine serum albumin (0.2 mg/ml). 5.2 μ l of the above solution was mixed with 1 μ g of λ gt 10 EcoR I arm, and then precipitated with ethanol. Recombinant phage DNA including both λ gt 10 and cDNA was prepared as follows. The above precipitate was reconstituted in 9 μ l of the column buffer and was incubated at 16 °C overnight by adding 1 μ l of 10mM adenosine triphosphate and 1 μ l of T4 DNA ligase (350 units/ μ l).

(4) Screening of cDNA library.

(i) Preparation of oligonucleotide probe

For preparation of probe, a mixture of complementary oligonucleotide of 17 mer (384 species mix) corresponding to the amino acid sequence from Val¹ to Pro⁶ in N-terminal amino acid sequence of TCF-II β -chain was synthesized and labelled 5' terminus with T4 polynucleotide kinase (TAKARA SHUZO) and [γ - ³²P]ATP (Amersham, Co, Ltd). This probe is shown as following;

complementary strand used as probe:
(384 species mix)

3' -CACCACCTTACCGTAGGG- 5'

G G G C A

A A A T

T T T

(ii) screening of recombinant phage

About 500 thousand plaques of phage were obtained by in vitro packaging of the recombinant phage DNA solution which was obtained by (3), using Gigapack Gold (Stratagene) and then by infecting to E.Coli C600hfl. After adsorption of the plaques to Hybond-N filter (Amersham), they were denatured with alkali, neutralized and baked at 80 °C for 2 hours. Hybridization was performed by the method of Bell et al (Nature 310 775 -777, 1984). The first screening was carried out by using the mix probe which was obtained from (1). One clone which would contain TCF-II fragment was found in the positive plaques detected by the first screening.

(5) Cloning of full length of TCF-II cDNA

Internal amino acid sequences (one letter code), (α) NYMGNLSQTRSGL and (β) TSXSVYGWGYTGILNYDGLL(X: not identified) were obtained respectively from α - and β -chains of TCF-II by digesting them with lysylendopeptidase and subsequently by mapping their fragments. N-terminal amino acid sequence of β -chain of TCF-II was coincided with that of β -chain of one of the human hepatocyte growth factors (hHGFs). Moreover, the above mentioned internal amino acid sequences (α) and (β) in TCF-II

were located in α - and β -chain of both hHGF, respectively. Therefore, it has been thought that TCF-II is expressed from one of the family of hHGF genes. MIYAAWA et al. (BBRC 163 ,967-973 (1989)) and NAKAMURA et al. (Nature. 342 ,440-443 (1989)) reported the hHGF cDNAs from placenta and liver cDNA libraries, respectively.

- 5 Comparison of all the primary sequence deduced from both hHGF cDNAs revealed differences in amino acids at 14 sites in their sequences. From these results, the presence of the family of hHGF genes was suggested. Identical regions between placenta type and liver type hHGF cDNAs were chosen as primer sequences for Polymerase Chain Reaction (PCR). The identical oligonucleotides of both hHGF cDNAs at 5' and 3' non-coding region were chemically synthesized and screening of TCF-II cDNA was carried out by
10 PCR using them as primers. Sal-77 primer which have a cleavage site of restriction enzyme Sal I and Sph2203 primer which have a cleavage site of restriction enzyme Sph I were synthesized by DNA synthesizer (Applied Co. Ltd). These primers are shown as following:

15 Sal-77 primer: 5'-GGTCGACTAGGCACTGACTCCGAACAGGATTC-3'

Sal I

20 Sph2203 primer: 5'-GGCATGCACAGTTGTATTGGTGGGTGCTTCAG-3'

Sph I

cloning with PCR method was carried out by following procedures.

- 25 (i) PCR

30	cDNA synthesized as described in (2) (dissolved in 150 μ l of column buffer)	1 μ l
	20 μ M Sal-77 primer	2.5 μ l
	20 μ M Sph2203 primer	2.5 μ l
	10 \times PCR reaction solution (100mM Tris HCl buffer (pH8.3) containing 500mM KCl, 15 mM MgCl ₂ and 0.1% (W/V) gelatin)	10 μ l
35	1.25mM dGTP, dATP, dTTP, dCTP, mixture	16 μ l
	Ampli Taq (5 units/ μ l TAKARASHUZO)	0.5 μ l
	Distilled water	67.5 μ l

- 40 After above solutions were mixed in microfuge tube with 0.5 ml volume size and covered on the liquid surface with 100 μ l of mineral oil (Sigma Co. Ltd). PCR was carried out by the Quick Thermo System (Japan Genetics Co. Ltd). After pretreatment at 94°C for 7 min, a three-step reaction which consists of annealing reaction, at 55°C for 3 min; polymerase reaction, at 72°C for 2min; and denature reaction, at 94°C for 2 min was repeated 35 times. Then the reaction mixture was treated for 3 min at 55°C , and
45 subsequently for 11min at 72 °C and then returned to room temperature (Each time includes the altering time of temperature). When a part of the reaction mixture was analyzed an electrophoresis using agarose gel, a DNA fragment consisting of about 2.3 kilobases (Kb) which was thought as aimed TCF-II cDNA, was obtained. Then, the DNA which was obtained from four tubes containing the above mentioned reaction mixture was precipitated with ethanol and digested with restriction enzymes, Sal I and Sph I . After an
50 agarose gel electrophoresis, DNA fragment with about 2.3Kb was recovered by using DE81 paper (Watman Co. Ltd).

- (ii) Subcloning

- 55 The DNA fragment with 2.3Kb digested with restriction enzymes, Sal I and Sph I , which was obtained from (1) , was inserted using ligation kit (TAKARA SHUZO) into a vector fragment which was onbtained by digestion of plasmid vector pUC18 (Japan gene Co. Ltd) with restriction enzymes, Sal I and Sph I , and transfected into Esherichia Coli DH5 α (according to protocol of BRL Co. Ltd). More than 20 subclones

could be obtained.

(iii) Determination of base sequence

5 The base sequences of obtained subclones were determined by the dideoxy-method (Sequenase Ver. 2.0 TOYOBO). Incorporation errors of nucleotides on the Ampi Taq (TAKARA SHUZO) were corrected by the analysis of base sequences of several subclones. The base sequence of TCF-II cDNA obtained by the above mentioned procedure and amino acid sequence deduced from the base sequence are shown in Fig. 15. It consists of 2172 base pairs (bp) from ATG of initiation codon for transcription to TAG of termination
10 codon. If translated into amino acid, TCF-II consists of 723 amino acids. Amino acid sequence from the first methionine (Met¹) to the 29 th alanine (Ala²⁹) residues is presumed as a signal sequence. As shown in Fig.15, TCF-II in which two polypeptides consisting of α chain and β chain are bound by disulfide bond is synthesized initially as a single chain. Since N-terminus of α chain in TCF-II was blocked, it was unidentified. However, N-terminal amino acid sequence of β chain and a few internal amino acid sequences of TCF-II
15 were determined as above mentioned, and were shown in Fig. 15.

The obtained base sequence of the TCF-II cDNA is very similar to hHGF which has been found by MIYAZAWA et al (Biochemical and Biophysical Research Communication 163 967-973 (1989)). TCF-II cDNA, however, deletes five amino acid residues (F-L-P-S-S) from Phe¹⁶² to Ser¹⁶⁶ in the amino acid sequence of hHGF. Therefore, the facts revealed that TCF-II cDNA was new one of the family of hHGF genes. The
20 comparison between the amino acid sequence of TCF-II, which is deduced from above mentioned base sequence, and the amino acid sequence of hHGF (MIYAZAWA et al) is shown in Fig. 16.

The method to obtain the new glycoprotein, TCF-II which is characterized by the above mentioned physicochemical properties is described here after.

Any human-derived fibroblast can be used as a cell for the production of the material in the present
25 invention. Human embryonic lung-derived fibroblast cells, human embryonic kidney-derived fibroblast cells, and human embryonic prepuce-derived fibroblast cells etc. are given as suitable cells. In the performance of the present invention, IMR-90 cells (ATCC CCL 186) and WI-38 cells (ATCC CCL 75) etc. are preferably suitable.

These cells are grown in serum-enriched medium or serum-free medium which are used in the
30 common cultures. Dulbecco's modified Eagle medium (DMEM) containing 5% bovine calf serum is given as the representative medium. Amino acids, transferin, fatty acids and hormones such as insulin etc. can be added if necessary.

The cells are cultured in the medium, and the standing culture using T flask etc., the floating culture using microcarrier, and the continuous culture using hollow fiber or ceramic carrier are able to be adopted.
35 It is preferable that the culture is carried out in atmosphere with 5% CO₂ at 20 to 37°C as the culture conditions and that the medium is exchanged every 2 to 3 days. After the cell density reaches the optimum, the medium is exchanged every 7 to 10 days and the culture broth is collected. The aimed glycoprotein is purified from the collected culture broth.

The collected culture broth is concentrated by UF concentration using a membrane with a pore size of
40 M.W. 6,000. The aimed glycoprotein in the UF concentrate is adsorbed to cation exchange resins and then eluted from the resins with buffer containing 0.3 ~ 0.6M NaCl. CM Sephadex (Pharmacia) C-50 etc. can be given as the ion exchange resins. The fractions which have the potent cytotoxic activity are collected and subsequently applied to affinity chromatography for glycoprotein. ConA-Sepharose is especially suitable to affinity chromatography for the aimed glycoprotein. The affinity column is equilibrated with 0.05M Tris-HCl
45 buffer, pH7.0, containing 0.5M NaCl, and then the above obtained fraction is applied to the column. After washing the column with the equilibration buffer, the active material is eluted from the column with an elution buffer containing carbohydrate corresponding to carbohydrate chain attached to the affinity column (or gel) for glycoprotein. When the above mentioned ConA-Sepharose is used, the active material is eluted with the buffer containing α -methyl-D-mannopyranoside. The eluted active fraction is dialysed against water
50 and lyophilized. The lyophilized active material is dissolved with 0.05M Tris-HCl buffer, pH6.0 to 7.0, containing 0.2M NaCl and is further purified on HPLC using a strong cation exchange column. Mono S (Pharmacia) is especially suitable to the strong cation exchange column. Elution of the active material from the Mono S column is carried out with a gradient from 0 to 1.0M NaCl and the active fractions are collected.

The present active material is eluted at salt intensity of 0.6 to 0.9M. Thus obtained active fraction is
55 further purified on an affinity chromatography using Heparin-Sepharose (Reg. TM) (Pharmacia). Elution of the active material from the Heparin-Sepharose (Reg. TM) column is carried out by a gradient from 0.3 to 2.0M NaCl and the aimed material is eluted at salt intensity of 1.0 to 1.5 M. Subsequently, assays for cytotoxic activity of the present inventional material, TCF-II to mouse L929-18 cells and for growth

stimulating activity of TCF-II for hepatocytes are mentioned below.

Assay for Cytotoxic Activity

5 Mouse L929 cells (ATCC CCL1) were subcloned and a subclone with the highest sensitivity to TCF-II in the present invention was selected. Thus, the clone, L-929-18 which had a high sensitivity to the tumor cell cytotoxic factor was obtained.

L-929-18 cells were grown to confluence in DMEM containing 10% FCS, and then the cells were harvested by trypsin treatment. The cells were suspended at a cell density of 6×10^5 cells/ml in DMEM containing 10% FCS and $1 \mu\text{g/ml}$ actinomycin D. $50 \mu\text{l}$ of DMEM which was prepared in the same way as the cell suspension was added to each well in 96-well microplate (Falcon) and $50 \mu\text{l}$ of the sample solution which contains the present substance, TCF-II, dissolved in the same DMEM was added to the first dilution well. Both were mixed well and $50 \mu\text{l}$ of the mixture was subsequently added to the second dilution well. A serially diluted substance was prepared by repeating the above procedures.

15 $50 \mu\text{l}$ of the cell suspension was inoculated into each well containing a serially diluted substance and the culture was carried out at 37°C for 2 days in a CO_2 incubator. After the culture, the medium was removed gently and the cells were washed twice with saline. The viable cells which adhere to each well were fixed and stained by addition of $50 \mu\text{l}$ of 0.5% crystal violet in the mixture of methanol and water (1:4) to each well. Each well was washed three times with distilled water and dried, and the crystal violet in each well was extracted with Sorenson's buffer (mixture of 6.1 ml of 0.1 M disodium citrate, 3.9 ml of 0.1N HCl, and 10 ml of ethanol). Absorbance of the extracts at 570 nm was determined by a microtiter spectrophotometer.

Unit of TCF-II ($\mu\text{g/ml}$) was defined as the dilution ratio given 50% cell death.

25 Assay for hepatocyte growth stimulating activity

Hepatocytes were separated from Wister male rat by the method of Segren (Method in cell biology, vol. 13, p29, Academic Press, New York, 1976). The obtained hepatocytes were planted on 24 well plastic plates (Falcon) at the cell density of 8.8×10^4 cells/0.5ml/well and cultured under the presence of 5% CO_2 at 37°C . A Williams E culture-medium (Flow Raboratory) which was supplemented with 10% fetal bovine serum (Hyclone), 100 u/ml penicillin and $100 \mu\text{g/ml}$ streptomycin was used as the culture medium (abbreviated below as basal culture medium). After incubation at 37°C for 24 hours, the culture medium was exchanged with the basal culture medium containing the test samples. The hepatocytes were further cultured for 24 hours and then cultured in the basal culture medium containing $4 \mu\text{Ci/ml}$ (86 Ci/m mol) of ^3H -thymidine (Amersham) for 2 hours followed by determination of DNA synthesis. When the hepatocytes were labeled with ^3H -thymidine, the incorporated counts (dpm) were determined from the difference of counts(dpm) which were measured between the presence and absence of 10 mM hydroxyurea on each test groups. After the cells were labeled by the above procedure, the cells were washed twice with cold PBS, 2% perchloric acid and 95% ethanol and dried by air, then dissolved in 0.8 ml of 2% SDS containing 2mM EDTA and 20mM NaHCO_3 followed by determination with a liquid scintillation counter.

The result is shown in Table 1.

Table 1

Sample	Concentration (ng/ml)	Hepatocyte growth stimulating activity (dpm/well, $\times 10_3$)
No addition	-	21.7 ± 9.2
hEGF	20	239.3 ± 7.2
TCF-II	1	93.7 ± 29.7
	10	378.5 ± 93.5
	100	467.4 ± 77.3

55 hEGF (WAKUNAGA Pharmacy Co Ltd) was used as the positive control for hepatocyte growth stimulating activity. The results in Table 1 demonstrate that hepatocyte growth stimulating activity of TCF-II is stronger than that of hEGF.

Nextly, the present invention describes pharmaceutical preparations which contains biologically active factor, TCF-II, as an active component.

The above mentioned biologically active factor in the present invention has following pharmaceutical activities:

① An antitumor activity:

TCF-II inhibits the growth of KB, HeLa, MCF-7 and BG-1 which is human-derived tumor cells, and have a cytotoxic activity to mouse L-929 cells, and to Sarcoma 180, Meth A Sarcoma and P388 of tumor cells, but not inhibit the growth of IMR-90 cells which are human-normal cells.

② A leukemia cell differentiation inducing activity:

TCF-II induces the differentiation of the human-derived leukemia cell line, HL-60 to granulocyte like cells.

③ A cellular immunology enhancing activity:

TCF-II stimulates the growth of human cytotoxic T cells

④ A vascular endothelial cell growth stimulating activity:

TCF-II stimulates the growth of human umbilical cord blood vessel-derived endothelial cells.

⑤ A hepatocyte growth stimulating activity:

TCF-II stimulates the growth of adult rat hepatocytes.

These activities are exhibited in very small dose range of 1-1000 ng/ml.

The pharmaceutical product in the present invention can be expected as an anti-tumor drug, an anti-leukemia drug, a cellular immunology enhancing drug, wound healing drug and therapeutic drug for liver disease including liver regenerating drug and so on. The biologically active factor (it is called as TCF-II hereafter) which is a glycoprotein with high molecular weight, however, is easily adsorbed to a polypropylene vessel and so on, which is an injection syringe, and to a glass vessel and so on.

TCF-II is also an unstable substance. The activity of TCF-II is considerably decreased by temperature or humidity and so on, and inactivated easily. Therefore, manufacturing of the stable pharmaceutical product is required. Accordingly, the pharmaceutical product in the present invention is a TCF-II one which contains one or more of proteins and nonionic detergents as adsorption preventors, or one or more of proteins, carbohydrates and aminoacids as stabilizing agents, or moreover the combination of one or more as adsorption preventors and one or more as stabilizing agents.

Any pharmaceutical form which is a lyophilized, or liquid or powder form can be used as a pharmaceutical product in the present invention, if the above mentioned adsorption preventors and stabilizing agents were included in the pharmaceutical product. The active component in the present invention, TCF-II can be produced or purified by the any method. TCF-II which is purified from culture broth of any TCF-II secreting cell, can be also used. TCF-II which is produced by the method of recombinant DNA technology, using Escherichia Coli, yeast, mammalian cells such as chinese hamster ovary etc, as hosts, and is purified by various methods, can be also used.

Albumin and gelatin etc, among proteins, and Tween 80 (Reg. TM) and Tween 20 (Reg. TM) etc, among nonionic detergents can be used as the adsorption preventor in the present invention.

Albumin and gelatin etc, among proteins; sorbitol, mannitol and xylitol etc, among carbohydrates; glycine and alanine etc, among amino acids can be used as the stabilizing agents in the present invention. The suitable added amounts of proteins as the adsorption preventors are more than 0.1%, preferably 0.1 to 20%, and the suitable added amounts of nonionic detergents are more than 0.001%, preferably 0.001 to 1.0%. The suitable added amounts of protein as stabilizing agents are more than 0.1%, preferably 0.1 to 20.0%; the suitable added amounts of carbohydrates are 5 to 40%; and the suitable added amounts of amino acids are more than 1%. On the case of using with combination of one or more of adsorption preventors and one or more of stabilizing reagents, or using with combination of two or more of stabilizing agents, the added amounts of each agent were permitted in the above mentioned range. The reasonable amounts of these adsorption preventors and stabilizing agents are dissolved and prepared as aqueous solution with suitable concentration and pH which can be used. The ratios of permeability pressure of this solutions are in the range of 0.1 to 3.1, preferentially 1.0. Since the stable range of TCF-II against pH is pH6 to 9, the aqueous solution for the preparation of pharmaceutical product is suitable to adjust pH to 6.0 to 9.0.

Then, the adsorption-prevention and stability of TCF-II in the pharmaceutical product in the present invention is further described in detail as shown in Experiments in the following:

Experiment 1. Test for prevention of adsorption

Human embryonic lung fibroblast, IMR-90 cells (ATCC, CCL186) were cultured for 7 days in DMEM containing 5% fetal bovine serum. 200 μ g of TCF-II which was purified from the culture broth was dissolved in 100 ml of 0.01M phosphate buffer, pH7.0, containing 0.15 M NaCl (PBS). Each 0.5ml of the TCF-II solution was added to glass tubes and polypropylene tubes. Separately, the solutions containing two-

fold, concentration of each added material shown in Table 1a, b and c were prepared with PBS. 0.5ml of the above solution with each concentration of added material was added to each tubes containing 0.5 ml of the TCF-II solution, and mixed well. The final concentration of TCF-II is 1 μ g/ml and the final concentration of each added material is adjusted to concentration described in Table 1 a, b and c.

5 As a control, 0.5ml of PBS was added to each glass tube or polypropylene tube containing 0.5 ml of the TCF-II solution.

Each test was carried out duplicately, and TCF-II activity was determined after the incubation for 1 hour at 37°C. Experimental values were shown as mean of duplicate incubations.

The activity of TCF-II was determined by cytotoxic activity as following.

10 A clone L929-18 with high sensitivity to TCF-II, which was obtained by subcloning mouse L929 (ATCC, CCL1), was used as a target cell.

L929-18 cells were grown to confluence in the Dulbecco's modified eagle medium (DMEM) containing 10% FCS, then harvested by trypsin treatment and by centrifugation. The obtained cells were suspended at a cell density of 6×10^5 cells/ml in DMEM containing 10% FCS and 1 μ g/ml of actinomycin D.

15 A serially diluted test sample was prepared by repeating dilution of test sample with DMEM which was used in the preparation of the cell suspension. Each 50 μ l of a serially diluted test sample was added to each well in 96 well-microplate (Falcon).

50 μ l of the cell suspension was inoculated into each well containing a serially diluted test sample and the culture was carried out at 37°C for 2 days in a CO₂ incubator. After the culture, the medium was removed gently and the cells were washed twice with PBS. The viable cells which adhere to each well were fixed and stained by addition of 50 μ l of 0.5% crystal violet in the mixture of methanol and water (1:4) to each well. Each well was washed with distilled water and dried, and crystal violet in each well was extracted with Sorenson's buffer (mixture of 6.1 ml of 0.1M disodium citrate, 3.9ml of 0.1N HCl, and 10ml of ethanol). Absorbances of the extracts at 570 nm were determined by a microtiter spectrophotometer.

25 Unit of TCF-II (u/ml) was defined as the dilution ratio given 50% cell death and the residual cytotoxic activity after the incubation was estimated as the relative activity(%) against the activity (100%) determined immediately after the preparation of test sample.

The results in Table 2,a, b, and c showed that TCF-II was easily adsorbed on surface of glass and polypropylene vessels and the adsorption preventors using in the present invention effectively prevented

30 TCF-II from the adsorption.

35

40

45

50

55

Table 2

(1) Glass tube

a. Effect of high molecular weight additives on
adsorption of TCF-II.

10

15

20

25

30

35

40

45

50

55

concentration	added material				
	human serum albumin(HSA)	low mol.w. gelatin*	gelatin	polyethylene glycol4000	dextran40
residual relative activity (%).....				
0	8.3	8.3	8.3	8.3	8.3
0.01	16.7	8.3	25.0	8.3	-
0.05	25.0	8.3	37.5	8.3	8.3
0.1	50.0	16.7	50.0	16.7	-
0.25	100.0	16.7	100.0	16.7	8.3
0.50	100.0	33.3	100.0	16.7	8.3
1.00	100.0	50.0	100.0	16.7	12.5
2.00	100.0	100.0	100.0	33.3	12.5
10.00	100.0	100.0	100.0	-	75.0
20.00	100.0	100.0	100.0	-	-

*: mean molecular weight = 6,000 (NIPPI Co. Ltd)

b. Effect of non-ionic detergents on adsorption of
TCF- II

5

10

added material

15

concentration
(%)

Tween80 (Reg.TM) Tween20 (Reg.TM)

.....residual relative activity (%).....

20

0

8.3

8.3

0.0001

16.7

16.7

0.0005

25.0

25.0

25

0.001

50.0

50.0

0.005

100.0

100.0

0.01

100.0

100.0

0.05

100.0

100.0

30

0.10

100.0

100.0

1.00

100.0

100.0

35

40

45

50

55

(2) Polypropylene tube

c. Effect of human serum albumin and Tween 80 (Reg. TM) on adsorption of TCF-II

5

10

15

20

25

30

35

added material		added material	
concentration	human serum albumin	concentration	Tween 80 (Reg. TM)
(%)	(HSA)	(%)	
residual relative activity		residual relative activity	
0	25.0	0	25.0
0.01	50.0	0.0001	33.3
0.10	75.0	0.0005	50.0
0.25	100.0	0.001	75.0
0.50	100.0	0.005	100.0
1.00	100.0	0.01	100.0
2.00	100.0	0.05	100.0
10.00	100.0	0.1	100.0
20.00	100.0	1.0	100.0

Experiment 2 Stability test

Under the conditions which prevented TCF-II from adsorption on surface of glass tube, effect of various kinds of added material on the stability of TCF-II was studied. 120 μ g of purified TCF-II from culture broth of IMR-90 cells was dissolved in 30 ml of PBS containing 0.02% Tween 80 (Reg. TM). After sterilization by filtrating the TCF-II solution using 0.22 μ filter, 0.5ml of the sterilized TCF-II solution was added to each sterilized glass tube.

The solution with two-fold concentration of each added material shown in Table 2 a, b and c, was prepared, and then sterilized by filtration with 0.22 μ filter. 0.5ml of the each solution was added to each glass tube containing 0.5ml of TCF-II solution and after mixing well, the glass tubes were sealed to prevent bacteria from contamination. As a control, 0.5 ml of PBS not containing Tween 80 (Reg. TM) was added to each glass tube containing 0.5 ml of TCF-II solution. The final concentration of TCF-II, Tween 80 (Reg. TM) and each added material were 2 μ g/ml, 0.01% and concentration shown in Table 3 a, b and c, respectively.

Each test was carried out duplicately. The TCF-II activity was determined after the incubation for one week at 40°C and estimated as relative activity (%) against the activity (100%) determined before the incubation. Experimental values were shown as mean of duplicate incubations.

From the results of Table 3 a, b and c, it was found that stabilizing agents which were used in the present invention had the effect which keeps on maintaining the activity of active component, TCF-II in the liquid state.

Table 3

a. Effect of high molecular weight additives on
stability of TCF-II *

10

		storage period (day) at 40°C			
	added		0	3	7
15	concentration				
	(%W/V)residual	relative	activity	(%).....
	human serum	0.0	100.0	25.0	16.7
	albumin	0.1	100.0	50.0	33.3
20		0.25	100.0	100.0	100.0
		0.5	100.0	100.0	100.0
	gelatin	0.0	100.0	25.0	16.7
		0.1	100.0	25.0	25.0
25		0.25	100.0	100.0	100.0
		0.5	100.0	100.0	100.0
	low mol.	0.0	100.0	25.0	16.7
30	weight	0.5	100.0	25.0	16.7
	gelatin	2.5	100.0	33.3	25.0
	(mean M.W. 6,000)				
35		0.0	100.0	25.0	16.7
	polyethylenglycol 4000				
		0.5	100.0	16.7	12.5
40		2.5	100.0	16.7	12.5

*:All test samples contained 0.01% Tween 80 (Reg. TM) to prevent TCF-II from adsorption
on surface of glass tube.

45

50

55

b. Effect of added carbohydrates on stability of TCF- II

5

		storage period (day) at 40°C		
10	added concentration (%W/V)	0	3	7
	residual relative activity (%).....		
15	Dextran40	0	100.0	25.0
		2	100.0	25.0
		10	100.0	12.5
20	Sorbitol	0	100.0	25.0
		2	100.0	33.3
		10	100.0	66.7
		20	100.0	100.0
		40	100.0	100.0
25	Mannitol	0	100.0	25.0
		2	100.0	33.3
		10	100.0	66.7
		20	100.0	100.0
		40	100.0	100.0
30	Glucose	2	100.0	16.7
		10	100.0	12.5
		20	100.0	25.0
		2	100.0	12.5
		10	100.0	12.5
35	Fructose	2	100.0	12.5
		10	100.0	<2.0

45

50

55

5	Mannose	0	100.0	25.0	16.7
		2	100.0	16.7	4.2
		10	100.0	<2.0	<2.0
10	Xylitol	0	100.0	25.0	16.7
		2	100.0	33.3	16.7
		10	100.0	100.0	66.7
		20	100.0	100.0	96.5

c. Effect of added amino acids on stability of TCF-II *

storage period (day) at 40°C					
amino acid	added concentration (%)	0	3	7residual relative activity (%).....
		0	3	7	
Arginine	0	100.0	25.0	16.7	
	1	100.0	25.0	16.7	
	5	100.0	50.0	33.3	
Glycine	0	100.0	25.0	16.7	
	1	100.0	33.3	25.0	
	5	100.0	100.0	66.7	
	10	100.0	100.0	100.0	
Lysine	0	100.0	25.0	16.7	
	1	100.0	25.0	16.7	
	5	100.0	66.7	16.7	
Alanine	0	100.0	25.0	16.7	
	1	100.0	25.0	16.7	
	5	100.0	100.0	50.0	
	10	100.0	100.0	90.0	

*:All test samples contained 0.01% Tween 80 (Reg. TM) to prevent TCF-II from adsorption on surface of glass tube.

Nextly, the results of pharmacological activities of TCF-II were shown in the following.

in vivo Antitumor Test to Sarcoma 180 of Human New Cytokine, TCF-II

Materials and methods

5 ① Experimental animals

Female ICR mice of 7 weeks old were purchased from Charles River Japan Inc.

10 ② Tumor cell lines

Sarcoma 180 cells were supplied by National Cancer Center, subcultured in mice with one time per week on this laboratory.

15 ③ Test samples

Test samples were prepared by dissolving TCF-II in 0.01M phosphate buffer, pH7.0 containing 0.8% NaCl, 0.01% Tween 80 (Reg. TM) and 0.25% human serum albumin.

Two series of test samples composed of 0.2 μ g TCF-II/0.2ml and 1.0 μ g TCF-II/0.2ml were prepared. To analyze the effect of pyrogen, test sample (940 pg pyrogen/0.2ml) containing standard pyrogen (Difco Inc.) corresponding to the amounts of pyrogen in 1 μ g of TCF-II was also prepared.

20 ④ Acute toxicity test

Acute toxicity test was carried out, using two mice group. Two doses, 10 μ g TCF-II/mouse and 20 μ g TCF-II/mouse were administered by injecting them into tail vein of mice, respectively. Toxicity was determined from mortality of animals.

25 ⑤ Antitumor test

Antitumor test was carried out, using seven mice/group.

Sarcoma 180 (10^6 cells/mouse) were planted under the skin of ICR mouse. The mice were divided in each test group and the test samples were injected into tail vein for 7 days at one time/every day. Inhibitory effect of tumor growth was determined by inhibitory ratio

$$35 \quad \frac{(C-T)}{C} \times 100\%$$

which was obtained from mean tumor weight (MTW) of injected groups against control group.

40 Results of tests

① Acute toxicity test

45 Toxicity was not appeared at doses of both 10 μ g and 20 μ g TCF-II/mouse.

50

55

② Antitumor test

Results after 3 weeks from the initiation of injection were shown in Table 4.

Table 4

Sample	Dose	MTW (mg)	$\frac{C-T}{C} \times 100(\%)$
control	0.0	3024.71	-
pyrogen	940 pg/mouse	3036.00	-0.37
TCF-II	0.2 μ g/mouse	1787.71	49.92
TCF-II	1.0 μ g/mouse	1984.21	34.40

Since on the above test, the optimum dose of TCF-II was unknown, test was not carried out by fitted doses. However, from these results, it was found that lower dose rather than high dose was much more effective.

Moreover, anti-tumor effect by direct injection to intratumor were also studied by the following method.

Methods:

Sarcoma 180 (1×10^6 cells/mouse) were planted under the skin of ICR mouse. The mice in which solid-tumors were vitally settled, were selected after one week of plantation. 0.2 μ g of TCF-II was injected for 7 days at one time/every day. When observation was continued for two weeks after the termination of injection, remarkable anti-tumor effect which caused necrosis of tumor site, changing into black was obtained. Moreover, mice in which tumor disappeared, were observed.

Brief Description of the Drawings

Fig.1 shows an elution profiles of TCF-II and plasminogen activator from the culture broth of IMR-90 cells containing 5% calf serum on CM-Sephadex C-50 chromatography. (1) and (2) shows the fractions eluted with 0.05M Tris-HCl buffer, pH7.0, containing 0.3M NaCl and with 0.05M Tris-HCl buffer, pH7.0, containing 0.6M NaCl, respectively.

-○-, -●- and -⊙- represent absorbance at 280 nm, plasminogen activator activity and cytotoxic activity to L929-18 cells, respectively.

Fig.2 shows the results of Con A affinity chromatography of TCF-II fraction which was eluted with 0.05M Tris-HCl buffer, pH 7.0, containing 0.6M NaCl on CM-Sephadex C-50 Chromatography of the culture broth of IMR-90 cells. (1) and (2) represent the washed fractions by 0.05M Tris-HCl buffer, pH7.0, containing 0.5M NaCl and the eluted fractions by 0.05M Tris HCl buffer, pH7.0, containing 0.5M NaCl and 0.3M α -methyl-D-mannopyranoside, respectively.

-●- and -⊙- represent absorbance at 280 nm and cytotoxic activity, respectively.

Fig. 3 shows an elution pattern on Mono S-HPLC of the TCF-II fraction which was obtained on Con A Sepharose affinity chromatography. -○- represents cytotoxic activity.

Fig.4 shows an elution profile of TCF-II on Heparin-sepharose affinity chromatography of the elute from Mono S-HPLC. (1) and (2) represent the washed fraction by 10mM Tris-HCl buffer, pH7.5, containing 0.3M NaCl and the eluted fraction by NaCl gradient from 0.3 to 2.0M, respectively. -●- and -⊙- represent absorbance at 280nm and cytotoxic activity, respectively.

Fig.5 shows SDS electrophoresis of TCF-II (reduction and non-reduction).

Fig.6 shows heat stability of TCF-II.

Fig.7 shows pH stability of TCF-II.

Fig.8 shows cytotoxic activity of TCF-II to human tumor cell lines in vitro.

Fig.9 shows cytotoxic activity of TCF-II to Sarcoma 180.

Fig.10 shows cytotoxic activity of TCF-II to Meth A and P388, respectively. -●- and -○- represent Meth A and P388, respectively.

Fig.11 shows cytostatic activity of TCF-II to human tumor cell lines. -●- and -○- represent cytostatic activities to ovarian carcinoma, BG-1 and breast carcinoma, MCF-7, respectively.

Fig. 12 and Fig.13 show effect of TCF-II on incorporation of ^3H -thymidine into the lymphocytes in the mixed lymphocytes culture at day 5 and at day 8, respectively. Six samples at day 5 and at day 8 were determined, respectively. Results are given as mean \pm SD.

Fig.14 shows the stimulatory effect of TCF-II on the growth of vascular endothelial cells, HUVEC.

Fig.15(1) and (2) show the base sequence of TCF-II cDNA and the amino acid sequence of TCF-II, which was deduced from the base sequence, respectively.

Fig.16 shows the comparison between the amino acid sequence of TCF-II, which was deduced from the above base sequence and an amino acid sequence of hHGF reported by Miyazawa et al.

A BEST FORM TO PUT ON THE INVENTION

The present invention is further described concretely as shown in EXAMPLES in the following:

EXAMPLE 1

(1) Culture of Human Fibroblast, IMR-90 cells

Human fibroblast, IMR-90 cells(ATCC CCL 186) (3×10^6 cells) were inoculated in one liter-roller bottle holding 100ml DMEM containing 5% calf serum (CS), and were cultured for 7 days with rolling at the rates of 0.5 to 2 rpm. When the total cell numbers reached 1×10^7 cells, the cells were harvested by trypsin treatment and collected in bottom of the bottle. After adding 250 ml of fresh DMEM containing 5% CS to the bottle, 100g of autoclaved ceramics with 5 to 9 mesh (Toshiba Ceramic. Co. Ltd) was added to the bottle containing cell suspension. Standing culture was carried out at 37°C for 24 hours. Then 250 ml of DMEM containing 5% CS was added to the bottle (final medium volume was 500 ml), and the culture was continued. Every 7 to 10 days all amount of the medium (about 500 ml) was collected and fresh medium (500ml) was supplied. Thus, the production was continued for 2 months and 4 l of the culture broth were collected per roller bottle. Specific activity of thus obtained culture broth was 32 u/ml.

(2)Purification of the Glycoprotein, TCF-II

75 liters of the culture broth described in (1) was concentrated about 10-fold by UF concentration, using Amikon-membrane filter (pore size, M.W. 6,000). Then, after CM-Sephadex C-50 was equilibrated with 0.05M Tris-HCl buffer, pH7.0, 1.5kg of wet weight of the resins was added to the above UF concentrated broth and the substance was absorbed to the resins by gently stirring at 4°C for 24 hour under pH 6.5 to 7.0. After the absorption, the resins were collected by filtrating through a Whatman No.2 filter paper. The collected resins were washed with 0.05M Tris-HCl buffer, pH7.0. Approximately 1500g of the washed resins was packed to the column (ϕ 7 x 40cm), and the column was eluted with 0.05M Tris-HCl, pH7.0, containing 0.01% Tween 20 (Reg. TM) and 0.3M NaCl. Elution of protein was monitored by absorbance at 280 nm. When protein was eluted almost completely, further elution at salt concentration of 0.6M NaCl was carried out. Cytotoxic activity to L929-18 cells and tissue plasminogen activator (t-PA) activity which IMR-90 cells produce were determined in each fraction. Thus obtained elution pattern is shown in Fig. 1. The fraction which was eluted at salt concentration of 0.6M NaCl exhibited the potent cytotoxic activity. This fraction was defined as TCF-II fraction. Then, Con A-Sepharose (Reg. TM) CL-6B (Pharmacia) was equilibrated with 0.05M Tris-HCl buffer, pH7.0, containing 0.5M NaCl, and the gel was packed into the column (ϕ 2.5 x 8cm). The column was washed well with the same buffer, the TCF-II fraction which was eluted from the CM-Sephadex column was loaded on the column. After the column was washed again with 0.05M Tris-HCl, pH7.0, containing 0.5M NaCl with 10 times volume of the column bed volume, the substance was eluted with 0.05M Tris-HCl, pH7.0, containing 0.5M NaCl and 0.3M α -methyl-D-mannopyranoside at a flow rate of 70 ml/hour. Elution of protein was monitored by optical absorbance at 280 nm and cytotoxic activity in each fraction was determined. Thus obtained elution pattern is shown in Fig.2. The fraction which was eluted firstly was collected and dialyzed against distilled water for 48 hours. The

dialyzed fraction was lyophilized and white powder was obtained. The lyophilized powder was dissolved with a small volume of 0.05M Tris-HCl buffer, pH7.0, containing 0.01% Tween 20 (Reg. TM) and 0.2M NaCl, and was loaded on MonoS column (Pharmacia) for HPLC which was equilibrated with 0.01M phosphate buffer, pH 7.0, containing 0.01% Tween 20 (Reg. TM). After the column was washed with 0.01M phosphate
5 buffer, pH7.0, containing 0.01% Tween 20 (Reg. TM) for 20 min at a flow rate of 0.5ml/min, elution was carried out by the NaCl gradient from 0 to 1.0M in 60min. The obtained elution pattern is shown in Fig.3. The active fraction was eluted at 0.76M NaCl. The active fraction was collected and diluted with the equilibration buffer and then loaded on the Mono S column for HPLC again. Elution was carried out again by the NaCl gradient from 0 to 1.0 M NaCl in the same buffer. Heparin-Sepharose (Reg. TM) (Pharmacia)
10 was equilibrated with 10 mM Tris buffer, pH 7.0, containing 0.3M NaCl, and 5ml of the gel was packed to the column (ϕ 1.0 \times 7cm). The active fraction from Mono S-HPLC was collected, diluted with 0.01M Tris-HCl buffer, pH7.0, to NaCl concentration of 0.3M, and loaded on the above mentioned column. Then, the column was washed with 0.01M Tris-HCl pH7.0, containing 0.3M NaCl with 10 times volumes of the column bed volume. The substance was eluted at a flow rate of 20 ml/hour by the NaCl gradient from 0.3M to 2.0
15 M in the same buffer. The elution pattern is shown in Fig.4. Thus, the purified glycoprotein was obtained. As shown in Table 5, 0.12 mg of the active glycoprotein was obtained from 75 liters of the culture broth as the starting material. This glycoprotein was a tumor cytotoxic factor and its specific activity was 5,248,000 u/mg.

20

25

30

35

40

45

50

55

Table 5

Purification of tumor cytotoxic factor obtained from the culture broth of iMR-90 (containing 5% CS)
TCF-II: A fraction eluted with 0.6M NaCl on Sephadex C-50 chromatography

Purification Step	Volume (ml)	Protein (mg/ml)	Total Protein (mg)	Cytotoxic Activity (u/ml)
Culture broth	75000	3.30	247500.0	32.0
UF Concentration	10000	23.40	234000.0	192.0
CM-Sephadex chromatography (eluted fraction with 0.6M NaCl)	894	0.32	205.6	2048.0
ConA Sepharose (Reg.TM) chromatography	244	0.26	63.4	5120.0
Mono S-HPLC	13	0.26	2.1	80000.0
Heparin-Sepharose (Reg.TM) chromatography	6	0.02	0.12	104960.0

Total Activity ($\times 10^4$ u)	Specific Activity (u.mg)	Purification Factor	Recovery (%)
24.0	9.7	1.0	100.0
192.0	8.2	0.9	80.0
183.1	8904.3	918.0	76.3
124.9	19692.3	2030.0	52.0
104.0	500000.0	51546.4	43.3
56.0	5248000.0	541030.9	23.3

Physicochemical properties of TCF-II obtained by the above mentioned procedure are shown as follows.

① Determination of molecular weight on SDS gel electrophoresis

Molecular weight of TCF-II was determined by electrophoresis using polyacrylamide gel containing 0.1% SDS. The glycoprotein showed two adjacent bands with M.W. 78,000 and 74,000. When the glycoprotein was reduced by 2-mercaptoethanol and the similar electrophoresis was carried out, three bands with molecular weight of 52,000, 32,000, and 28,000 were observed (Fig.5).

These results indicate that TCF-II is a heterodimer composed of a common subunit with M.W. 52,000 and a subunit of M.W. 28,000 or a subunit of M.W. 32,000.

② Isoelectric point

Isoelectric point of the substance was determined as 7.4 to 8.55 by isoelectric focussing, using Phast Gel IEF3-9.

③ Heat stability

TCF-II which had the activity of 51,200 u/ml was added to 0.1M Tris-HCl buffer, pH7.0, containing 0.01% Tween 20 and the TCF-II solution with 512 u/ml was prepared. The solution was treated for 10 min at 25, 35, 50, 60, 70, 80, 90, and 95 °C. The residual cytotoxic activity (u/ml) after treatment at each temperature was estimated as the relative activity (%) against the activity (u/ml, 100%) at 25 °C (control). As shown in Fig.6, the substance was stable till 60 °C.

④ pH stability

Each buffer containing 0.01% Tween 20 (Reg. TM), composition of which was shown in Table 6, was prepared. The amount of TCF-II corresponding to 51,200 u/ml when it was prepared at pH 8 was dissolved in each buffer, and allowed to stand at 37 °C for 1 hour. The relative activity (%) compared to the control at room temperature for one hour was determined. As shown in Fig. 7, the glycoprotein was stable in the range of pH 6 to 9.

Table 6

Preparation of buffers	
pH1-3	1/10 M Glycine-HCl
pH4-6	1/10 M Acetate buffer
pH7-8	1/10 M Tris-HCl
pH9.12	1/10 M Glycine-NaOH

⑤ N-terminal amino acid sequence of TCF-II.

50 µg of TCF-II was reduced and three polypeptides, A with M.W. 52,000, B with M.W.32,000, and C with M.W. 28,000, were separated by the electroblot method. Amino acid sequence of each polypeptide was analyzed using the Applied Biosystems 477A Protein Sequencer. N-terminal amino acid sequence of polypeptide A could not be determined because its N-terminus had been blocked. Polypeptides B and C had the common N-terminal amino acid sequence as follows;

Val-Val-Asn-Gly-Ile-Pro-Thr-X-Thr-Asn-Ile-Gly-

1 2 3 4 5 6 7 8 9 10 11 12

X-Met-Val-Ser-Leu-

13 14 15 16 17

X means an unidentified amino acid.

Since polypeptide B and C show the same N-terminal amino acid sequence, TCF-II appears to have dimer structure in which polypeptide A with M.W. 52,000 is bound to polypeptide B with M.W. 32,000 or to polypeptide C with M.W. 28,000 by S-S bridge.

⑥ Amino acid composition

Ten μ g of TCF-II, which was determined by BioRad Protein Assay kit, was hydrolyzed with HCl and then its amino acid composition was determined using a Hitachi Amino Acid analyzer, Model L-8500.

Amino acid composition of the material is shown as follows.

Amino acid composition:		
A.A	nmol	mol%
Asp	10.375	12.97
Glu	7.750	9.69
Ser	5.000	6.25
Gly	7.250	9.06
His	3.000	3.75
Arg	5.375	6.72
Thr	5.125	6.41
Ala	2.625	3.28
Pro	5.625	7.03
Tyr	3.875	4.84
Val	4.125	5.16
Met	1.875	2.34
Cys	ND	-
Ile	5.000	6.25
Leu	4.875	6.09
Phe	2.250	2.81
Trp	ND	-
Lys	5.875	7.34
total	80.000	100(99.99)

EXPERIMENT 2

Cytotoxic activity to the tumor cells of the glycoprotein, TCF-II obtained in EXPERIMENT 1 is shown.

① Inhibition of the growth of tumor cells

The human tumor cell lines, HeLa and KB, and the human normal diploid cells, IMR-90 were suspended at a cell density of 1×10^5 cells/ml in DMEM containing 10% FCS, respectively. 50 μ l of each cell suspension was added to each well in a 96 well microplate (Falcon). Each 50 μ l of 10-, 20-, 40-, 80- and 160- fold dilution of TCF-II solution (5,120 u/ml) with DMEM was added to each well containing each cell suspension, and the mixture was cultured at 37°C for 3 days in a CO₂ incubator. The surviving cells in each well were fixed and stained by addition of 50 μ l of 0.5% crystal violet solution in the mixture of methanol and water (1:4) to each well. Each well was washed with distilled water and dried, and the crystal violet in each well was extracted with Serenson's buffer. Absorbances of the extracts at 570nm were determined by a microtiter spectrophotometer.

Cell growth inhibition (%) of TCF-II was calculated compared to the group in the absence of TCF-II as the control, and plotted against the concentration of TCF-II. As shown in Fig. 8, TCF-II showed the potent inhibition of the growth of KB and HeLa cells whereas it did not inhibit the growth of normal cells, IMR-90.

② Reaction with antibodies for the known materials

TCF-II was dissolved in DMEM containing 10% FCS to the concentration of 320 u/ml. The titer of anti-LT antibody which neutralized 1000 u/ml LT was added to the TCF-II solution and the mixture was allowed to stand at 37°C for 1 hour. Similarly, anti-TNF antibody and anti-INF- β were added to the TCF-II solution at the concentrations of 1×10^6 u/ml and 1000 u/ml, respectively. Each antibody used in this experiment is commercially available.

After the reactions, cytotoxic activity of TCF-II was determined. Any antibody, however, did not neutralize the activity.

EXPERIMENT 3

Cytotoxic activity to various mouse tumor cell lines of the glycoprotein, TCF-II obtained in EXPERIMENT 1 is shown.

Sarcoma 180, Meth A sarcoma and P-388 cells were used as mouse tumor cell lines.

Sarcoma 180 cells were suspended in DMEM containing 10% fetal bovine serum and Meth A and P-388 cells were suspended at 2×10^4 cells/ml, in RPMI 1640 medium containing 10% fetal bovine serum, respectively. 50 μ l of each cell suspension was inoculated into each well in 96-well microplate (Falcon). TCF-II was dissolved in DMEM containing 10% fetal bovine serum for Sarcoma 180 cells and in RPMI 1640 medium containing 10% fetal bovine serum for Meth A and P-388 cells. Each 50 μ l of TCF-II solution which was prepared so that the final concentrations of TCF-II were 0, 2, 4, 8, 16, 31, 62, 125, 250, 500 and 1000 ng/ml, respectively, was added to each well containing each cell suspension. After each cell line were cultured at 37°C for 3 days in a CO₂ incubator. The cells in each well were stained with Trypan blue and the viable cell number were counted using a hemacytometer. The viable cell numbers were expressed as mean of duplicate experiments. Cytotoxic activity (%) was calculated according to the following equation.

Cytotoxic activity(%) =

$$\frac{\text{Average surviving numbers in control (cells/ml)} - \text{Average surviving cell numbers in TCF-II group (cells/ml)}}{\text{Average surviving cell numbers in control (cells/ml)}} \times 100$$

Cytotoxic activities of TCF-II to Sarcoma 180 cells and to Meth A sarcoma and P-388 cells were shown in Fig.9 and Fig.10, respectively.

All cell lines were highly sensitive to TCF-II, and the IC₅₀ values for cytotoxic activity of TCF-II on Sarcoma 180, Meth A and P-388 cells were 6, 40 and 460 ng/ml, respectively.

EXPERIMENT 4

Inhibitory effect of the glycoprotein, TCF-II obtained in Experiment 1 on human tumor cell lines, ovarian carcinoma, BG-1 and breast carcinoma, MCF-7 is shown in the following.

BG-1 and MCF-7 cells were suspended in McCoy medium containing 10% FCS and in Eagle's MEM containing 10% FCS, non-essential amino acid mixture, pyruvate and Eagle's salts at 2×10^4 cells/ml, respectively. TCF-II was dissolved in McCoy medium containing 10% FCS for BG-1 cells and in Eagle's MEM containing 10% FCS for MCF-7 and a serially diluted TCF-II solution was prepared by repeating 2-fold dilution of 4 μ g TCF-II/ml with the same medium.

50 μ l of each cell suspension was inoculated into each well in 96-well microplate (Falcon). Each 50 μ l of a serially diluted solution of TCF-II which was prepared for each cell line was added to the wells containing each cell suspension. The culture was carried out at 37°C for 5 days in CO₂ incubator. After the culture, the culture broth was removed and the cells were gently washed twice with PBS. The surviving cells which adhered to each well were fixed and stained by addition of 50 μ l of 0.5% crystal violet solution in the mixture of methanol and water (1:4) to each well. Each well was washed with distilled water and dried, and crystal violet in each well was extracted with Serenson's buffer. Absorbances of the extracts at 570 nm were determined by a microtiter spectrophotometer. Cell growth inhibition (%) was calculated for each cell line according to the following equation compared to the control in the absence of TCF-II.

Cell growth inhibition(%) =

$$\frac{\text{Control OD}_{570\text{nm}} - \text{TCF-II group OD}_{570\text{nm}}}{\text{Control OD}_{570\text{nm}}} \times 100$$

Results are shown in Fig.11. These results indicate that TCF-II inhibits the growth of both tumor cell lines, BG-1 and MCF-7.

EXPERIMENT 5

Promyelocytic leukemia cell line, HL-60 cells -differentiation inducing activity of the glycoprotein, TCF-II obtained in EXPERIMENT 1 is shown.

HL-60 cells were suspended at a cell density of 3.5×10^5 cells/ml in RPMI 1640 medium containing 10% fetal bovine serum. 100 μ l of the cell suspension was added to each well in 96-well flat bottomed microplate(Falcon). Then, 100 μ l of the TCF-II solution in the same medium was added to each well containing cell suspension to give the final concentrations of 15.6, 62.5, 125, 250, 500, and 1000 ng/ml.

The cells were cultured at 37°C for 3 and 7 days, and HL-6 cells-differentiation inducing activity of TCF-II was determined by nitroblue tetrazolium (NBT) reduction assay. Moreover, morphological change of the cells was observed.

1)NBT reducing ability

NBT reducing ability is shown in Table 7.

Table 7

30

35

40

TCF-II concentration (ng/ml)	NBT reducing ability(%)	
	Culture days	
	3	7
0	7.4	11.7
15.6	10.6	20.4
62.5	11.1	24.5
125	12.4	28.3
250	16.9	45.2
500	12.4	29.6
1000	12.1	26.8

The values shown in Table 7 represent percentage of cells containing blue-black formazan deposits (average of 2 experiments) when at least more than 200 cells were counted.

(When HL-60 cells differentiate to normal cells, they get NBT reducing ability and accumulate blue-black formazan in them)

This result has demonstrated that TCF-II induces the differentiation of promyelocytic leukemia cell line, HL-60 and has the highest activity for the differentiation induction at 250ng/ml.

2)Morphological change

It is known that HL-60 differentiates to two way of macrophages and granulocytes by differentiation inducing factors.

Changes in morphology and nucleus of the cells which were cultured with TCF-II at 37°C for 7 days were investigated by the light Giemsa staining method. The result indicates that HL-60 differentiates to granulocyte-like cells.

EXPERIMENT 6

Stimulatory effect on cellular immunology of the glycoprotein, TCF-II obtained in EXPERIMENT 1 is shown.

The effect of TCF-II on blastgenic transformation of lymphocytes was observed when mixed lymphocyte culture was carried out in the presence of TCF-II.

Lymphocytes were isolated from human peripheral blood by the method of Ficall-Conray and suspended in RPMI 1640 containing 10% FCS. Lymphocytes from two persons were mixed as 1:1 and added to wells in 96 well round bottomed microplate at 1×10^5 cells/ 100 μ l/well. TCF-II was added at various concentrations and the cells were cultured in RPMI 1640 containing 10% FCS in a CO₂ incubator. ³H-Thymidine was added at 0.25 μ Ci/well at 16 hours before the end of culture. After the culture the cells were harvested by a cell harvester and washed with PBS. The radioactivity of ³H-Thymidine incorporated into the cells were determined by a scintillation counter.

Results are given in Fig.12 and Fig.13. TCF-II did not show the stimulatory effect at day 5 of the culture as shown in Fig.12, but ³H-thymidine incorporation was significantly stimulated by the presence of TCF-II compared to the control at day 8 as shown in Fig.13. These results indicate that TCF-II stimulates the growth of cytotoxic T cell, that is, TCF-II possesses the enhancing effect on cellular immunology.

EXPERIMENT 7

Stimulatory effect on vascular endothelial cell growth of the glycoprotein, TCF-II obtained in EXPERIMENT 1 is shown.

Human umbilical vein endothelial cells, HUVEC was used as a test cell. The endothelial cells, HUVEC, were suspended at a cell density of 2.5×10^4 cells/ml in E-GM medium containing 2% fetal bovine serum. 50 μ l of the cell suspension was added to each well in 96-well flat bottomed microplate (Falcon). Then, 50 μ l of the TCF-II solution which was prepared in the same medium was added to give the final concentrations of 0, 4, 8, 16, 31, 62, 125, 250, 500 and 1000 ng/ml. The cells were cultured at 37°C for 6 days in a CO₂ incubator. After the culture, the culture medium in each well was removed and the cells were washed gently with PBS. The cells were removed from the well by trypsin treatment and viable cell numbers were counted using a heamacytometer.

The effect of TCF-II on human normal vascular endothelial cells is shown in Fig.14. The result indicates that TCF-II does not show the cytotoxic activity to the normal cells but has the stimulatory effect on the growth. Especially, the cell growth stimulatory activity was the maximal at a concentration of 125 μ g/ml.

The following Experiments show prescriptions of the pharmaceutical product in the present invention.

EXPERIMENT 8

TCF-II	20 μ g
Human serum albumin	100mg

The above mentioned compositions were dissolved in 0.01M phosphate buffer, pH7.0, containing 0.15M NaCl (PBS) and the resulted solution was filled up to 20ml with PBS. After sterilization by filtrating the obtained solution using 0.22 μ filter, 2ml of the sterilized solution was poured into each vial tube and lyophilized, and then the vial tubes were sealed.

EXPERIMENT 9

TCF-II	40 μ g
Tween80 (Reg.TM)	1mg
Human serum albumin	50mg

The above mentioned compositions were dissolved in physiological salt solution (0.8%NaCl) and the solution was filled up to 20 ml with the same solvent. After the sterilization by filtrating the solution using 0.22 μ filter, 2ml of the sterilized solution was poured into each vial tube and lyophilized, and then the vial

tubes were sealed.

EXPERIMENT 10

5

TCF-II	20 μ g
Tween80 (Reg.TM)	2mg
Sorbitol	4g

10

The above mentioned compositions were dissolved in PBS, and the resulted solution was filled up to 20 ml with PBS. After sterilization by filtrating the solution using 0.22 μ filter, 2ml of the sterilized solution was poured into each vial tube and lyophilized, and then the vial tubes were sealed.

EXPERIMENT 11

20

TCF-II	40 μ g
Tween80 (Reg.TM)	2mg
Glycine	2g

25

The above mentioned compositions were dissolved in physiological salt solution (0.8%NaCl), and the resulted solution was filled up to 20 ml with the same solvent. After sterilization by filtrating the solution using 0.22 μ filter, 2ml of the sterilized solution was poured into each vial tube and lyophilized, and then the vial tubes were sealed.

EXPERIMENT 12

30

TCF-II	40 μ g
Tween80 (Reg.TM)	1mg
Sorbitol	2g
Glycine	1g

35

The above mentioned compositions were dissolved in physiological salt solution (0.8%NaCl), and the resulted solution was filled up to 20 ml with the same solvent. After sterilization by filtrating the solution using 0.22 μ filter, 2ml of the sterilized solution was poured into each vial tube and lyophilized, and then the vial tubes were sealed.

EXPERIMENT 13

45

TCF-II	20 μ g
Sorbitol	4g
Human serum albumin	50mg

50

The above mentioned compositions were dissolved in PBS and the resulted solution was filled up to 20 ml with PBS. After sterilization by filtrating the solution using 0.22 μ filter, 2ml of the sterilized solution was poured into each vial tube and lyophilized, and then the vial tubes were sealed.

55

EXPERIMENT 14

5

TCF-II	40 μ g
Glycine	2g
Human serum albumin	50mg

10 The above mentioned compositions were dissolved in physiological salt solution (0.8%NaCl), and the resulted solution was filled up to 20 ml with the same solvent. After sterilization by filtrating the solution using 0.22 μ filter, 2ml of the sterilized solution was poured into each vial tube and lyophilized, and then the vial tubes were sealed.

Industrial Availability

15

The present invention provides a new glycoprotein. The glycoprotein in the present invention can be used as a tumor cytotoxic factor, leukemia cell line-differentiation inducing factor, cellular immunology enhancing factor and vascular endothelial cell growth factor, and can be provided usually.

20 Moreover, the glycoprotein in the present invention can be used as a biochemical or pharmacological reagent.

Claims

1. A glycoprotein which (i) is obtainable from the culture broth of human derived fibroblasts, (ii) has amino acid sequence shown in Fig. 15 and (iii) has the following physicochemical properties;
 - a. Molecular weight; On the determinations of molecular weight by SDS polyacrylamide gel electrophoresis, 78,000 \pm 2,000 or 74,000 \pm 2,000 under the nonreduced conditions and a common band A with 52,000 \pm 2,000 and band B with 30,000 \pm 2,000 or band C with 26,000 \pm 2,000 under the reduced conditions.
 - b. Isoelectric point; 7.4 to 8.6.
 - c. Heat stability; Stable in heating at 60 °C for 10min.
 - d. pH stability; Stable in the range of pH6 to 9.
 - e. Carbohydrate chain; Adsorbed to a Concanavalin A (Con A)-Sepharose (Reg. TM) column.
 - f. Biological activity; Inhibits the growth of KB cells, Hela cells, and L-929 cells but not IMR-90 cells.
 - g. Reactivity to antibodies; The cytotoxic activity is not neutralized by anti-TNF antibody, anti-lymphotoxin antibody, and anti-interferon- β antibody.
2. A pharmaceutical product, which contains the glycoprotein according to claim 1 as active component, moreover one or more of proteins and nonionic detergents as adsorption preventor, and/or one or more of proteins, carbohydrates and amino acids as stabilizing agent, and which is characterized by the following biological activity
 - ① An antitumor activity:
Inhibits the growth of KB, HeLa, MCF-7 and BG-1 which are human-derived tumor cell lines, and have a cytotoxic activity to mouse L-929 cells, and to Sarcoma 180, Meth A sarcoma and P388 which are mouse tumor cell lines, but not inhibit the growth of IMR-90 cells which are human-normal cells.
 - ② A leukemia cell differentiation inducing activity:
Induces the differentiation of the human-derived leukemia cell line, HL-60 to granulocyte-like cells.
 - ③ A cellular immunology enhancing activity:
Stimulates the growth of human cytotoxic T cells
 - ④ A vascular endothelial cell growth stimulating activity:
Stimulates the growth of human umbilical vein endothelial cells, HUVEC.
 - ⑤ A hepatocyte growth stimulating activity:
Stimulates the growth of adult rat hepatocytes.
3. A pharmaceutical product of biological factor according to claim (2), in which the protein selected as the adsorption preventor is either albumin or gelatin.

4. A pharmaceutical product of biological factor according to claim (2), in which the nonionic detergent selected as the adsorption preventor is either Tween 80 (Reg. TM) or Tween 20 (Reg. TM).
5. A pharmaceutical product of biological factor according to claim (2), in which the protein selected as the stabilizing agent of the pharmaceutical product is either albumin or gelatin.
6. A pharmaceutical product of biological factor according to claim (2), in which the carbohydrate selected as the stabilizing agent of the pharmaceutical product is sorbitol, mannitol or xylitol.
7. A pharmaceutical product of biological factor according to claim (2), in which the amino acid selected as the stabilizing agent of the pharmaceutical product is either glycine or alanine.
8. A pharmaceutical product of biological factor according to claim (2), which contains various combination of one or more of proteins or nonionic detergents as adsorption preventor, and one or more of proteins, carbohydrates or amino acids as stabilizing agent.
9. A DNA containing base sequence encoding amino acid sequence showed in Figure 15.
10. A cDNA containing base sequence encoding amino acid sequence showed in Figure 15.
11. A cDNA sequence containing base sequence showed in Figure 15.

Patentansprüche

1. Glycoprotein,
 - (i) das erhältlich ist aus der Kulturbede von auf den Menschen zurückgehenden Fibroblasten,
 - (ii) eine Aminosäuresequenz gemäß Figur 15 besitzt und
 - (iii) die folgenden physikochemischen Eigenschaften besitzt:
 - (a) Molekulargewicht: bei Bestimmung des Molekulargewichts durch SDS-Polyacrylamid-Gel-Elektrophorese 78.000 ± 2.000 oder 74.000 ± 2.000 unter nicht-reduzierenden Bedingungen mit einer gemeinsamen Bande A bei 52.000 ± 2.000 und einer Bande B bei 30.000 ± 2.000 oder einer Bande C bei 26.000 ± 2.000 unter reduzierenden Bedingungen.
 - (b) Isoelektrischer Punkt: 7,4 bis 8,6;
 - (c) Wärmestabilität: Stabil beim Erhitzen bei 60°C für 10 min;
 - (d) pH-Stabilität: Stabil im Bereich von pH 6 bis 9;
 - (e) Kohlenwasserstoffkette: Adsorbiert an eine Concanavalin A-Sepharose^R-Säule (Con-A-Sepharose^R-Säule);
 - (f) biologische Aktivität: Inhibiert das Wachstum von KB-Zellen, Hela-Zellen und L-929-Zellen, jedoch nicht von IMR-90-Zellen;
 - (g) Reaktivität gegenüber Antikörpern: Die cytotoxische Aktivität wird nicht durch Anti-TNF-Antikörper, Anti-Lymphotoxin-Antikörper und Anti-Interferon- β -Antikörper neutralisiert.
2. Pharmazeutisches Produkt, das das Glycoprotein gemäß Anspruch 1 als Wirkstoff, ferner ein oder mehrere Proteine und nichtionische Detergenzien als Adsorptions Inhibitoren und/oder ein oder mehrere Proteine, Kohlenhydrate und Aminosäuren als Stabilisatoren enthält und durch die folgende biologische Aktivität gekennzeichnet ist:
 - (1) Antitumor-Aktivität: Inhibiert das Wachstum von KB, HeLa, MCF-7 und BG-1, bei denen es sich um vom Menschen abgeleitete Tumorzelllinien handelt und die eine cytotoxische Aktivität gegenüber Maus-L-929-Zellen und gegenüber Sarcoma-180, Meth-A-Sarcoma und P388 besitzen, die Maustumorzelllinien sind, inhibiert jedoch nicht das Wachstum von IMR-90-Zellen, bei denen es sich um normale Humanzellen handelt;
 - (2) Aktivität zum Induzieren einer Leukämie-Zellen-Differenzierung: Induziert die Differenzierung einer vom Menschen abgeleiteten Leukämie-Zelllinie HL-60 zu granulocytenartigen Zellen;
 - (3) Aktivität zur Beschleunigung zellulärer Immunologie: Stimuliert das Wachstum von cytotoxischen Human-T-Zellen;
 - (4) Aktivität zum Stimulieren eines vaskulären Endothelial-Zellen-Wachstums: Stimuliert das Wachstum von Human-Umbilicalvenen-Endothelialzellen HUVEC;

(5) Aktivität zum Stimulieren von Hepatocytenwachstum: Stimuliert das Wachstum von adulten Rattenhepatocyten.

3. Pharmazeutisches Produkt nach Anspruch 2, bei dem das Protein, das als Adsorptionsinhibitor gewählt wurde, entweder Albumin oder Gelatine ist.
4. Pharmazeutisches Produkt nach Anspruch 2, bei dem das nichtionische Detergens, das als Adsorptionsinhibitor gewählt ist, entweder Tween 80^R oder Tween 20^R ist.
5. Pharmazeutisches Produkt nach Anspruch 2, bei dem das Protein, das als Stabilisator für das pharmazeutische Produkt gewählt ist, entweder Albumin oder Gelatine ist.
6. Pharmazeutisches Produkt nach Anspruch 2, bei dem das Kohlenhydrat, das als Stabilisator für das pharmazeutische Produkt gewählt ist, Sorbit, Mannit oder Xylit ist.
7. Pharmazeutisches Produkt nach Anspruch 2, bei dem die Aminosäure, die als Stabilisator für das pharmazeutische Produkt gewählt ist, entweder Glycin oder Alanin ist.
8. Pharmazeutisches Produkt nach Anspruch 2, enthaltend verschiedene Kombinationen von einem oder mehreren Proteinen oder nicht-ionischen Detergenzien als Adsorptionsinhibitoren und ein oder mehrere Proteine, Kohlenhydrate oder Aminosäuren als Stabilisatoren.
9. DNA, enthaltend eine Basensequenz, die die Aminosäuresequenz gemäß Figur 15 kodiert.
10. cDNA, enthaltend eine Basensequenz, die eine Aminosäuresequenz gemäß Figur 15 kodiert.
11. cDNA-Sequenz, enthaltend eine Basensequenz gemäß Figur 15.

Revendications

1. Glycoprotéine qui (i) peut être obtenue à partir du bouillon de culture de fibroblastes d'origine humaine, (ii) a la séquence d'acides aminés présentée dans la figure 15 et (iii) a les propriétés physico-chimiques suivantes:
 - a. Poids moléculaire: d'après les déterminations de poids moléculaire par électrophorèse sur gel de SDS-polyacrylamide, 78 000 \pm 2000 ou 74 000 \pm 2000 dans les conditions non réduites et une bande commune A à 52 000 \pm 2000 et une bande B à 30 000 \pm 2000 ou une bande C à 26 000 \pm 2000 dans les conditions réduites.
 - b. Point isoélectrique: 7,4 à 8,6.
 - c. Stabilité à la chaleur: stable au chauffage à 60 °C pendant 10 minutes.
 - d. Stabilité au pH: stable dans un intervalle de pH 6 à 9.
 - e. Chaîne hydrate de carbone: adsorbée à une colonne de concanavaleine A (Con A)-Sephrose (marque déposée).
 - f. Activité biologique: inhibe la croissance des cellules KB, des cellules HeLa et des cellules L-929, mais non des cellules IMR-90.
 - g. Réactivité vis-à-vis des anticorps: l'activité cytotoxique n'est pas neutralisée par l'anticorps anti-TNF, l'anticorps anti-lymphotoxine ni l'anticorps anti-interféron β .
2. Produit pharmaceutique qui contient la glycoprotéine selon la revendication 1 comme composant actif, ainsi qu'une ou plusieurs protéines et/ou détergents non ioniques comme agents empêchant l'adsorption, et un ou plusieurs protéines, hydrates de carbone et acides aminés comme agents stabilisateurs, et qui se caractérise par l'activité biologique suivante:
 - (1) une activité antitumorale: inhibe la croissance de KB, HeLa, MCF-7 et BG-1 qui sont des lignées cellulaires tumorales d'origine humaine, et ont une activité cytotoxique vis-à-vis des cellules L-929 de souris, et de Sarcome 180, Sarcome Meth A et P388 qui sont des lignées cellulaires tumorales de souris, mais n'inhibent pas la croissance des cellules IMR-90 qui sont des cellules humaines normales.
 - (2) une activité d'induction de la différenciation des cellules leucémiques: induit la différenciation de la lignée cellulaire de leucémie d'origine humaine HL-60 en cellules de type granulocytes.

- (3) une activité d'accroissement de l'immunité cellulaire: stimule la croissance des cellules T cytotoxiques humaines.
- (4) une activité de stimulation de la croissance des cellules endothéliales vasculaires: stimule la croissance des cellules endothéliales de la veine ombilicale humaine, HUVEC.
- 5 (5) une activité de stimulation de la croissance des hépatocytes: stimule la croissance des hépatocytes de rat adultes.
3. Produit pharmaceutique du facteur biologique selon la revendication 2, dans lequel la protéine choisie comme agent empêchant l'adsorption est l'albumine ou la gélatine.
- 10 4. Produit pharmaceutique du facteur biologique selon la revendication 2, dans lequel le détergent non ionique choisi comme agent empêchant l'adsorption est le Tween 80 (marque déposée) ou le Tween 20 (marque déposée).
- 15 5. Produit pharmaceutique du facteur biologique selon la revendication 2, dans lequel la protéine choisie comme agent stabilisant du produit pharmaceutique est l'albumine ou la gélatine.
- 20 6. Produit pharmaceutique du facteur biologique selon la revendication 2, dans lequel l'hydrate de carbone choisi comme agent stabilisateur du produit pharmaceutique est le sorbitol, le mannitol ou le xylitol.
7. Produit pharmaceutique du facteur biologique selon la revendication 2, dans lequel l'acide aminé choisi comme agent stabilisateur du produit pharmaceutique est la glycine ou l'alanine.
- 25 8. Produit pharmaceutique du facteur biologique selon la revendication 2, qui contient diverses combinaisons d'un(e) ou plusieurs protéines ou détergents non ioniques comme agents empêchant l'adsorption, et une ou plusieurs protéines, hydrates de carbone ou acides aminés comme agent stabilisateurs.
- 30 9. ADN contenant une séquence de bases codant pour la séquence d'acides aminés présentée dans la figure 15.
10. ADNc contenant la séquence de bases codant pour la séquence d'acides aminés présentée dans la figure 15.
- 35 11. Séquence d'ADNc contenant la séquence de bases présentée dans la figure 15.


40

45

50

55

FIG. 1







FIG. 2







FIG. 3









FIG. 4





F I G . 5



F I G . 6




FIG. 7






FIG. 8





F I G . 9





FIG. 10







FIG. 11






FIG. 14




Fig. 15 (1)

-110 -100 -90 -80 -70 -60
 TAGGCACTGACTCCGAA
 -50 -40 -30 -20 -10 0
 CAGGATTCTTTTACCCAGGCATCTCCTCCAGAGGGATCCGCCAGCCCGTCCAGCAGCACC
 10 20 30 40 50 60
 ATGTGGGTGACCAAACTCCTGCCAGCCCTGCTGCTGCAGCATGTCTCCTGCATCTCCTC
 M W V T K L L P A L L L Q H V L L H L L
 70 80 90 100 110 120
 CTGCTCCCATCGCCATCCCTATGCAGAGGGACAAAGGAAAAAGAAATACAATTCAT
 L L P I A I P Y A E G Q R K R R N T I H
 130 140 150 160 170 180
 GAATTCAAAAATCAGCAAAGACTACCCTAATCAAAATAGATCCAGCACTGAAGATAAA
 E F K K S A K T T L I E I D P A L K I E
 190 200 210 220 230 240
 ACCAAAAAGTGAATACTGCAGACCAATGTGCTAATAGATGTACTAGGAATAAAGGACTT
 T K K V N T A D Q C A N R C T R N K G L
 250 260 270 280 290 300
 CCATTCACCTTGCAAGGCTTTTGTGTTTGAATAAGCAAGAAACAATGCCTCTGGTTCCCC
 P F T C K A P V F D K A R K Q C L W F P
 310 320 330 340 350 360
 TTCAATAGCATGTCAAGTGGAGTGAATAAGAAATTTGGCCATGAATTTGACCTCTATGAA
 F N S M S S G V K K E F G H E F D L Y E
 370 380 390 400 410 420
 AACAAAGACTACATTAGAACTGCATCATTGGTAAAGGACGCAGCTACAAGGGAACAGTA
 N K D Y I R N C I I G K G R S Y K G T V
 430 440 450 460 470 480
 TCTATCACTAAGAGTGGCATCAAATGTGAGCCCTGGAGTTCATGATACCACACGAACAC
 S I T K S G I K C Q P W S S N I P H E H
 490 500 510 520 530 540
 AGCTATCGGGTAAAGACCTACAGGAAACTACTGTGAAATCCTCGAGGGGAAGAAGGG
 S Y R G K D L Q E N Y C R N P R G E E G
 550 560 570 580 590 600
 GGACCCTGGTGTTCACAAGCAATCCAGAGGTACGCTACGAAGTCTGTGACATTCCTCAG
 G P W C F T S N P E V R Y E Y C D I P Q
 610 620 630 640 650 660
 TGTTCAAGAAGTTGAATGCATGACCTGCAATGGGGAGAGTTATCGAGGTCTCATGGATCAT
 C S E Y E C M T C N G E S Y R G L M D H
 670 680 690 700 710 720
 ACAGAATCAGGCAAGATTTGTGACCGCTGGGATCATCAGACACCACACCGGCACAAATTC
 T E S G K I C Q R V D H Q T P H R H K F
 730 740 750 760 770 780
 TTGCCTGAAAGATATCCCGACAAGGGCTTTGATGATAATTATTGCCGCAATCCCGATGGC
 L P E R Y P D K G F D D N Y C R N P D G
 790 800 810 820 830 840
 CAGCCGAGGCCATGGTGTATACTCTTGACCCCTCACACCCGCTGGGAGTACTGTGCAATT
 Q P R P W C Y T L D P H T R V E Y C A I
 850 860 870 880 890 900
 AAAACATGCGCTGACAATACTATGAATGACACTGATGTTCTTTGGAACAACCTGAATGC
 K T C A D N T M N D T D V P L E T T E C
 910 920 930 940 950 960
 ATCCAAGCTCAAGGAGAAGGCTACAGGGGCACTGTCAATACCATTGGAATGGAATTCCA
 I Q G Q G E G Y R G T V N T I V N G I P
 970 980 990 1000 1010 1020
 TGTGAGCGTTGGGATTCTCAGTATCCTCAGGAGCATGACATGACTCCTGAAAAATTTCAAG
 C Q K W D S Q Y P H E H D M T P E N F K
 1030 1040 1050 1060 1070 1080
 TGCAAGGACCTACGACAAAATTAAGTCCGAAATCCAGATGGGTCTGAATCACCCTGGTGT
 C K D L R E N Y C R N P D G S E S P W C
 1090 1100 1110 1120 1130 1140
 TTTACCACTGATCCAAACATCCGAGTTGGCTACTGCTCCCAATTCAAAATCTGTGATATG
 F T T D P N I R V G Y C S Q I P N C D M




Fig. 15 (2)

1150 1160 1170 1180 1190 1200
 TCACATGGACAAGATTGTTATCGCTCGGAATGGCAAAATTATATGGGGCAACTTATGCCAA
 S H G Q D C Y R G N G K N Y M G N L S Q
 internal amino acid sequence
 1210 1220 1230 1240 1250 1260
ACAAGATCTGGACTAACATGTTCAATGTGGCACAACAACATGGAAGACTTACATCGTCAT
 T R S G L T C S M V D K N M E D L H R M
 in α -chain
 1270 1280 1290 1300 1310 1320
 ATCTTCTGGGAACCAGATGCAAGTAAGCTCAATGAGAATTACTGCCGAAATCCAGATGAT
 I F V E P D A S K L N E N Y C R N P D D
 1330 1340 1350 1360 1370 1380
 GATCGTCATGGACCCCTGGTGGCTACACGGGAAATCCACTCATTCCCTGGGATTATTGCCCT
 D A H G P V C Y T G N P L I P V W D Y C P
 1390 1400 1410 1420 1430 1440
 ATTTCTCGTTCTGAAGGTGATACCAACCTACAATACTCAATTTAGACCATCCCGTAATA
 I S R C E G D T T P T I Y N L D M P V I
 1450 1460 1470 1480 1490 1500
 TCTTGTCCCAAAACGAAACAAATTCGGAGTTGTAAATGGGATTCCAAACAGCAACAAACATA
 S C A K T K Q L H Y V N G I P T R T N I
 N-terminal in β -chain N-terminal amino acid sequence
 1510 1520 1530 1540 1550 1560
GGATGGATGGTTAGTTTGAGATACAGAAATAACATATGTCCGGAGGATCATTGATAAAG
 G V M Y S L R Y R N K H I C G G S L I K
 in β -chain
 1570 1580 1590 1600 1610 1620
 CAGACTTGGCTTCTTACTGCACGACAGTGTTCCTTCTCGAGACTTGAAAGATTATGAA
 E S W Y L T A R Q C F P S K D L K D Y E
 1630 1640 1650 1660 1670 1680
 GCTTGGCTTGGAAATTCATGATGTCCACGGGAACAGGAGATGAGAAATGCCAAACAGGTTCTC
 A W L G I H D V H G R G D E K C K Q Y L
 1690 1700 1710 1720 1730 1740
 AATGTTTCCACGCTGGTATATGCCCTGAAGGATCAGATCTGGTTTAAATGAAGCTTGCC
 N Y S Q L Y Y G P E G S D L Y L M K L A
 1750 1760 1770 1780 1790 1800
 AGGCCTGCTGCTCGCATGATTTTGTAGTACGATTGATTTACCTAATTATCGATGCACA
 R P A Y L D D F V S T I D L P N Y G C T
 1810 1820 1830 1840 1850 1860
ATTCTGAAAGACACGATTCGAGTGTTTATGGCTGGGCTACACTGGATTGATCAACTAT
 I P E K T S C S Y Y G V G Y T G L I N Y
 internal amino acid sequence
 1870 1880 1890 1900 1910 1920
GATGGCTATTACCAAGTGGCACATCTCTATATAATGGGAAATGAGAAATGCAGCCAGCAT
 D G L L R Y A H L Y I M G N E K C S Q H
 in β -chain
 1930 1940 1950 1960 1970 1980
 CATCGAGGGAAGGTGACTCTGAATCAGTCTGAAATATGTGCTGGGGCTGAAAAGATTGGA
 H R G K V T L N E S E I C A G A E K I G
 1990 2000 2010 2020 2030 2040
 TCAGGACCATGTGAGGGGGATTATGGTGGCCCACTTGTGTTGTGAGCAACATAAAATGAGA
 S G P C E G D Y G G P L Y C E Q H K M R
 2050 2060 2070 2080 2090 2100
 ATGGTTCTTGGTGTCAATTGCTGGTGGATGTGCCATTCCAAATCGTCCTGCTATT
 M Y L G V I Y P G R C C A I P N R P G I
 2110 2120 2130 2140 2150 2160
 TTTGTCCGAGTAGCATATTATGCAAAATGGATACACAAAATTATTTAATCATATAAGCTA
 F Y R Y A Y Y A K W I H K I I L T Y K Y
 2170 2180 2190 2200 2210
 CCACAGTCATAGCTGAAGTAAGTGTGTCTGAAGCACCCACCAATACAAGTGT
 P Q S A




Fig. 16

```

1
MWVTKLLPALLLQHIVLLHLLLLPIAIPYAEGQRKRNTIHEFKKSAKTTLIKIDPALKIK
*****
MWVTKLLPALLLQHIVLLHLLLLPIAIPYAEGQRKRNTIHEFKKSAKTTLIKIDPALKIK
1

61
TKKYNTADQCANRCTRNKGLPFTCKAFYFDKARKQCLWFPFNSMSSGVKKEFGHEFDLYE
*****
TKKYNTADQCANRCTRNKGLPFTCKAFYFDKARKQCLWFPFNSMSSGVKKEFGHEFDLYE
61

121
NKDYIRNCIGKGRSYKGTYSITKSGIKCQPWSSMIPHEH-----SYRGKDLQENYCRNP
*****
NKDYIRNCIGKGRSYKGTYSITKSGIKCQPWSSMIPHEHSFLPSSYRGKDLQENYCRNP
121

176
RGEEGGPWCFTSNPEVRYEYCDIPQCSEVECMTNCGESYRGLMDHTESGKICQRWDHQTP
*****
RGEEGGPWCFTSNPEVRYEYCDIPQCSEVECMTNCGESYRGLMDHTESGKICQRWDHQTP
181

236
HRHKFLPERYPDKGFDDNYCRNPDGQPRPWCYTLDPHTRWEYCAIKTCADNTMNDTDVPL
*****
HRHKFLPERYPDKGFDDNYCRNPDGQPRPWCYTLDPHTRWEYCAIKTCADNTMNDTDVPL
241

296
ETTECIQQQGEgyrgtYNTIVNGIPCQRWDSQYPHEHDMTPENFKCKDLRENYCRNPDGS
*****
ETTECIQQQGEgyrgtYNTIVNGIPCQRWDSQYPHEHDMTPENFKCKDLRENYCRNPDGS
301

356
ESPWCFTTDPNIRYGYCSQIPNCDMSHGQDCYRGNGXNYMGNLSQTRSGLTCSMWDKNME
*****
ESPWCFTTDPNIRYGYCSQIPNCDMSHGQDCYRGNGXNYMGNLSQTRSGLTCSMWDKNME
361

416
DLHRHIFWEPDASKLNYCRNPDDAHGPWCYTGNPLIPWDYCPISRCEGDTTPTIVNL
*****
DLHRHIFWEPDASKLNYCRNPDDAHGPWCYTGNPLIPWDYCPISRCEGDTTPTIVNL
421

476
DHPVISCATKQLRVVNGIPTRTNIGWVYSLRYRNKHICGGSLIKESWVLTARQCFFSRD
*****
DHPVISCATKQLRVVNGIPTRTNIGWVYSLRYRNKHICGGSLIKESWVLTARQCFFSRD
481

536
LKDYEAWLGIHDVHGRGDEKCKQVLNYSQLYYGPEGSDLYLMLKLARPAVLDDPVSTIDL
*****
LKDYEAWLGIHDVHGRGDEKCKQVLNYSQLYYGPEGSDLYLMLKLARPAVLDDPVSTIDL
541

596
NYGCTIPEKTSYVYGWYTGlinYDGLLRVAHLYIMGNEKCSQHHRGKVTLNESEICAG
*****
NYGCTIPEKTSYVYGWYTGlinYDGLLRVAHLYIMGNEKCSQHHRGKVTLNESEICAG
601

656
AEKIGSGPCEGDYGGPLYCEQHHRMVLGVIVPGRGCAIPNRPFI FVRVAYYAKVIHKII
*****
AEKIGSGPCEGDYGGPLYCEQHHRMVLGVIVPGRGCAIPNRPFI FVRVAYYAKVIHKII
661

716      723
LTYKVPQS
*****
LTYKVPQS
721      728

```

